

The orbifold fundamental group of Persson-Noether-Horikawa surfaces.

Fabrizio Catanese and Sandro Manfredini

This article is dedicated to the
memory of Boris Moishezon.

1 Introduction.

Among the minimal surfaces of general type, the Noether surfaces are those for which the Noether inequality $K^2 \geq 2p_g - 4$ is an equality (K^2 is the self intersection of a canonical divisor, p_g is the dimension of the space of holomorphic 2-forms).

These surfaces were described by Noether ([No]) and more recently by Horikawa ([Ho]) who proved that if $8 \mid K^2$ then there are two distinct deformation types, namely the Noether-Horikawa surfaces of connected type (for short, N-H surfaces of type C), and those of non connected type (for short, of type N). This notation refers to the fact that, the canonical map being a double covering of a rational ruled surface, for type C the branch locus is connected, whereas for type N it is not connected.

In particular Horikawa proved that the intersection forms are both of the same parity (in fact, both odd) if and only if $16 \mid K^2$.

From M. Freedman's theorem ([Fr]) follows that if $16 \mid K^2$ type N and type C provide two orientedly homeomorphic compact 4-manifolds.

Horikawa posed the question whether type N and type C provide two orientedly diffeomorphic compact 4-manifolds.

It looked like a natural problem to try to see whether the two differentiable structures could be distinguished by means of the invariants introduced by S. Donaldson in [Do].

In the case of type C we have been able ([Ca]) to calculate the constant Donaldson invariants (corresponding to zero-dimensional moduli spaces) using some singular canonical models of these surfaces with very many singularities, and an approach introduced by P. Kronheimer ([Kr]) for the case of the Kummer surfaces. The number we obtained, namely 2^{2k} when $K^2 = 8k$, is the leading term of the Donaldson series (see [K-M]), which was later fully calculated by Fintushel and Stern in the case of N-H surfaces of type C via the technique of

rational blow-downs ([F-S]).

The Donaldson series for N-H surfaces of type N has not yet, to our knowledge, been calculated; although, after the Seiberg-Witten theory ([W]) has been introduced, and after Pidstrigach and Tyurin ([P-T]) have announced the equality between Kronheimer-Mrowka and Seiberg-Witten classes, the two series should be equal.

Our original aim was to extend the application of the Kronheimer theory to the case of N-H surfaces of type N using a very singular model constructed by Ulf Persson ([Per]), describing its orbifold fundamental group, its representations into $SO(3)$, and then trying to see which of those have virtual dimension zero. In this article we consider the singular N-H surfaces of type N with maximal Picard number constructed by Persson, henceforth called Persson-Noether-Horikawa surfaces (P-N-H for short), and we determine their orbifold fundamental group.

This is our main result:

Theorem. *The orbifold fundamental group of the P-N-H surfaces is*

$$\mathbb{Z}_4 \oplus \mathbb{Z}_2$$

if $16 \mid K^2$,

$$\mathbb{Z}_4 \oplus \mathbb{Z}_4$$

in the other case where $8 \mid K^2$ but 16 does not divide K^2 .

It follows immediately that we have, for $16 \mid K^2$, only six nontrivial classes of orbifold $SO(3)$ -representations, and a result which we do not prove here is that we do not get anyone of virtual dimension zero.

This is not surprising in view of ([P-T]), since if Kronheimer's approach would have worked, we would have had only a finite number of constant Donaldson invariants.

On the other hand, the algebro-geometric technique of studying canonical models with many rational double points produces on the smooth model configurations of (-2) -projective lines (spheres) whose tubular neighborhood has a unique holomorphic structure and, in particular, a unique compatible C^∞ structure. In this way one produces a decomposition of the 4-manifold in geometric pieces, one of which is the nonsingular part of the singular canonical model.

From this point of view, the calculation of the orbifold fundamental group leads to a better understanding of the differentiable structures of the smooth model. Since our proof is rather involved technically we would like to give a brief geometrical "explanation" of our result.

Persson's construction starts with a plane nodal cubic C meeting a conic Q at only one point P . Moreover, C and Q have two common tangents L_{-1} and L_1 which meet in a point O collinear with P and the node of C .

Blowing up O we get a $\mathbb{P}^1$ -bundle $f' : \mathbb{F}_1 \rightarrow \mathbb{P}^1$ with a section Σ_∞ , a bisection Q' and a 3-section C' (' denoting the proper transform under the blow up).

A cyclic cover of order $2k+2$ branched on L'_{-1} and L'_1 yields a new $\mathbb{P}^1$ -bundle

$f'' : \mathbb{F}_{2k+2} \rightarrow \mathbb{P}^1$ with a section Σ''_∞ disjoint from a 3-section C'' and two sections Q''_1, Q''_2 (the inverse image Q'' of Q' splits into two components).

The curve $B = C'' \cup Q''_1 \cup Q''_2 \cup \Sigma''_\infty$ has many singular points, and our canonical model X_{2k+2} is the double cover of $\mathbb{F}_{2k+2}$ branched on B . By construction X_{2k+2} has a genus 2 fibration onto $\mathbb{P}^1$, whence the orbifold fundamental group $\pi_1(X_{2k+2}^\#)$, $X_{2k+2}^\#$ being the nonsingular part of X_{2k+2} , is a quotient of $\pi_1(F)$, where F is a fixed genus 2 fibre.

F being a double cover of $\mathbb{P}^1$ branched in six points $P_0 = \mathbb{P}^1 \cap \Sigma''_\infty$, $P_1 = \mathbb{P}^1 \cap Q''_1$, $P_2 = \mathbb{P}^1 \cap Q''_2$, $\{P_3, P_4, P_5\} = \mathbb{P}^1 \cap C''$, $\pi_1(F)$ is the subgroup of a free product $\mathcal{F}_5(2)$ of five copies of $\mathbb{Z}_2$, given by words of even length.

$\mathcal{F}_5(2)$ is generated by elements $\varepsilon_1, \dots, \varepsilon_6$ such that $\varepsilon_1 \cdots \varepsilon_6 = 1$ (ε_i corresponds to a loop in $\mathbb{P}^1$ around the point P_{i-1}).

The first main point (we must be rather vague here, else we must give the full proof) is that, since curve C'' is irreducible, when the fibre F moves around, $\varepsilon_4, \varepsilon_5, \varepsilon_6$ become identified.

Thus we only have $\varepsilon_1, \dots, \varepsilon_4$ with $\varepsilon_1 \cdots \varepsilon_4 = 1$, and therefore we have "proved" that our group is abelian, being a quotient of the fundamental group Γ of a curve of genus 1 obtained as the double cover of $\mathbb{P}^1$ branched in four points. More precisely, Γ is an abelian group with generators $\varepsilon_1 \varepsilon_2, \varepsilon_1 \varepsilon_3$.

We must still take into account the fact that, when the fibre F moves towards a singular point (corresponding to points of intersection $C'' \cap Q''_1, C'' \cap Q''_2, Q''_1 \cap Q''_2$), further relations are introduced. These relations are hard to control globally but if we look locally around these points of intersection, and accordingly take a new basis $\varepsilon'_1, \dots, \varepsilon'_4$, the situation becomes simpler.

In fact, the local equation of the double cover is $z^2 = y^2 - x^{2c}$, where $c=6$ or $c=k+1$, and x is the pullback of a local coordinate on $\mathbb{P}^1$, so that the corresponding local braid yields the relation $(\varepsilon'_j \varepsilon'_i)^c = (\varepsilon'_i \varepsilon'_j)^c$. In turn, using $(\varepsilon'_i)^2 = 1$, we obtain the relation $(\varepsilon'_j \varepsilon'_i)^{2c} = 1$.

That's how one shows that the two generators of the abelian group have period 2 or 4.

The paper is organized as follows:

In section two we take up Persson's construction using explicit equations showing that the surface is defined over a real quadratic field.

In the third section we describe the five steps leading to a presentation of our fundamental group in terms of the braid monodromy of the plane curve $D = C \cup Q$. Finally, in section four we apply combinatorial group theory arguments in order to give the main result concerning the orbifold fundamental group.

Acknowledgements : Both authors acknowledge support from the AGE Project H.C.M. contract ERBCHRXCT 940557 and from 40% M.U.R.S.T..

The first author would like to express his gratitude to the Max-Planck Institut in Bonn where this research was initiated (in 1993), and to the Accademia dei Lincei where he is currently Professore Distaccato.

2 Persson's configuration.

In this section we will provide explicit equations for the configuration constructed by Ulf Persson in [Per].

This is the configuration formed by a smooth conic Q and a nodal cubic C intersecting in only one point P which is smooth for C . Moreover Q and C have two common tangents L_1 and L_{-1} meeting in a point O lying on the line joining P and the node of C .

Let $Q \subset \mathbb{C} \mathbb{P}^2$ be the conic $\{(x, y, z) \in \mathbb{C} \mathbb{P}^2 \mid x^2 + 2zy + z^2 = 0\}$.

Since

$$x^2 + 2zy + z^2 = (x+z)^2 + 2z(y-x) = (x-z)^2 + 2z(y+x)$$

Q is tangent to the lines $L_1 = \{x-y=0\}$ and $L_{-1} = \{x+y=0\}$.

The tangency points are:

$$x-y=x+z=0 \Rightarrow (1, 1, -1)$$

$$x+y=x-z=0 \Rightarrow (1, -1, 1).$$

Note that Q is also tangent to the line $z=0$ at the point $(0, 1, 0) = P$.

We want to find an irreducible nodal cubic C such that $C \cdot Q = 6P$ and such that C is tangent to the lines $x = \pm y$ in points different from those of Q .

Let C be a cubic s.t. $P \in C$ and $C \cdot Q = 6P$. Note that if C were reducible, then the previous condition would imply that $z=0$ is a component of C .

We then have $\text{div}(C) = \text{div}(z^3) \pmod{Q}$, so $C = z^3 + QL$ with L a linear form, and thus

$$C = z^3 + (x^2 + 2zy + z^2)(ax + by + cz).$$

Since we want C to be tangent to the two lines L_1 and L_{-1} we obtain that the following homogeneous polynomials in (x, z)

$$z^3 + (x+z)^2((a+b)(x+z) + z(c-a-b)) \tag{2.1}$$

$$z^3 + (x-z)^2((a-b)(x-z) + z(c+a-b)) \tag{2.2}$$

must have a double root.

Set $\zeta = (\frac{z}{x+z})$ and $\hat{\zeta} = (\frac{z}{x-z})$ and rewrite 2.1, 2.2 as:

$$\zeta^3 + \zeta(c-a-b) + (a+b) = 0 \quad \hat{\zeta}^3 + \hat{\zeta}(c+a-b) + (a-b) = 0.$$

We recall that if ζ is a double root of $z^3 + pz + q = 0$ then

$$3\zeta^2 + p = 0 \quad \text{whence} \quad \frac{2}{3}\zeta p + q = 0$$

and this implies that

$$\zeta = -\frac{3}{2} \frac{p}{q} \quad \text{thus} \quad 27q^2 + 4p^3 = 0.$$

Therefore we have a double root of 2.1 if and only if

$$\exists A : \zeta=A, q=2A^3, p=-3A^2, \text{ i.e. } \begin{cases} a+b=2A^3 \\ c-(a+b)=-3A^2. \end{cases}$$

Similarly if we set $\hat{\zeta}=-B$ we have

$$\begin{cases} a-b=-2B^3 \\ c-(b-a)=-3B^2 \end{cases}$$

and so

$$\begin{cases} a=A^3-B^3 \\ b=A^3+B^3 \\ c=2A^3-3A^2=2B^3-3B^2. \end{cases}$$

Then A and B must satisfy $2(A^3-B^3)=3(A^2-B^2)$.

Recall that (we make no distinction between a curve and its equation)

$$C = z^3 + (x^2 + 2zy + z^2)((A^3 - B^3)x + (A^3 + B^3)y + (2A^3 - 3A^2)z)$$

while $x-y=0$ is tangent to C at the point where

$$\zeta = \frac{z}{x+z} = A.$$

Therefore the tangency point is $(1-A, 1-A, A)$.

Similarly $x+y=0$ is tangent to C at the point $(B-1, 1-B, B)$.

Let us now search for a cubic C with a singular point on the line $x=0$, as in Persson's construction.

Since $\frac{\partial C}{\partial x}$ on the line $x=0$ equals aQ and the singular point is different from P it follows that $a=0$. Whence $A^3-B^3=A^2-B^2=0$ and so $A=B$.

If $A=B$ then C contains only the monomial x^2 as a polynomial in x , so the involution $x \mapsto -x$ leaves the curve C invariant. From this we deduce that a singular point of C must have its x coordinate equal to 0 and C has then a singularity on the line $x=0$ if and only if

$$z^3 + (z^2 + 2zy)(2A^3y + (2A^3 - 3A^2)z) \text{ has a double root.}$$

Remembering that it can't be $A=B=0$, the double root cannot be $z=0$ and we can write the above as

$$z(z^2 + (z+2y)(2A^3y + (2A^3 - 3A^2)z)).$$

So we must check that

$$\begin{aligned} & z^2(1+2A^3-3A^2) + 2zy(A^3+2A^3-3A^2) + 4A^3y^2 = \\ & = z^2(1+2A^3-3A^2) + 2zy3A^2(A-1) + 4A^3y^2 \end{aligned}$$

has a double root.
This is the case when

$$9A^4(A-1)^2=4A^3(1+2A^3-3A^2) \quad \text{i.e.}$$

$$9A^6-18A^5+9A^4=4A^3+8A^6-12A^5.$$

Upon dividing by $A^3 \neq 0$ we get

$$A^3-6A^2+9A-4=0.$$

Observe that 1 is a root of this equation, but if $A=1$ then the singular point is $(0,0,1)$ and coincides with the point of tangency of $x+y=0$ so this root has to be discarded. Since

$$A^3-6A^2+9A-4=(A-1)(A^2-5A+4)=(A-1)^2(A-4)$$

the other possible root is then $A=4$, and in this case we have $B=A=4$, $a=0$, $b=8 \cdot 4^2$, $c=5 \cdot 4^2$.

Then

$$C = z^3 + 4^2(x^2 + 2yz + z^2)(8y + 5z)$$

The tangency points are $(-3, -3, 4)$ and $(3, -3, 4)$, while for the singular point we have $x=0$ and a double root of

$$z^2 + 4^2(2y + z)(8y + 5z) = 0 \iff 81z^2 + 4^2 18zy + 4^4 y = 0 \iff 9z + 4^2 y = 0$$

so the singular point is $(0, 9, -16)$.

With this choice of A and B , C is irreducible (since $z=0$ is not a component of C).

We want to find the lines through $(0, 0, 1)$ and tangent to C .

Let $A=B=\lambda$ and consider more generally the 1-parameter family of curves:

$$C_\lambda = z^3 + (x^2 + 2yz + z^2)(2\lambda^3 y + (2\lambda^3 - 3\lambda^2)z) = 0.$$

The tangency points on the two fixed lines $x+y=0$, $x-y=0$ are, as we know, $(1-\lambda, 1-\lambda, \lambda)$ and $(\lambda-1, 1-\lambda, \lambda)$.

Rewriting the last equation in powers of z we obtain:

$$z^3(1+2\lambda^3-3\lambda^2) + z^2 6y\lambda^2(\lambda-1) + z\lambda^2(4\lambda y^2 + (2\lambda-3)x^2) + 2\lambda^3 x^2 y = 0.$$

Since we know what happens for $\lambda=0$, we can divide by λ^3 , set $w = \frac{z}{\lambda}$ and obtain:

$$w^3(1+2\lambda^3-3\lambda^2) + w^2 6y\lambda(\lambda-1) + w(4\lambda y^2 + (2\lambda-3)x^2) + 2x^2 y = 0.$$

We let now Δ be the discriminant of C_λ with respect to the variable w , and using a standard formula for Δ , we find a degree 6 equation in x and y which

is divisible by $x^2(x^2-y^2)$.

Remembering that the discriminant of $a_0x^3+a_1x^2+a_2x+a_3$ is:

$$\Delta = a_1^2a_2^2 - 4a_0a_2^3 - 4a_1^3a_3 - 27a_0^2a_3^2 + 18a_0a_1a_2a_3$$

and applying this formula for simplicity when $\lambda=4$, we obtain:

$$y^22^63^4(16y^2+5x^2)^2 - 2^23^4(16y^2+5x^2)^3 - \\ - 2^{12}3^6x^2y^4 - 2^23^{11}x^4y^2 + 2^53^8(16y^2+5x^2)x^2y^2$$

and factoring this binary form we get:

$$x^2(x^2-y^2)2^23^4(2^7y^2-5^3x^2).$$

So we have that the tangent lines to C passing through $(0,0,1)$ are $x = \pm y$, $x = \pm \sqrt{\frac{128}{125}}y$ while $x=0$ passes through the node of C . We denote by L_0 the line

$x=0$ and by L_+, L_- the two lines $x = \sqrt{\frac{128}{125}}y$, $x = -\sqrt{\frac{128}{125}}y$ respectively.

In order to find the tangency point on the lines L_+, L_- we by symmetry may restrict to the line L_+ .

Writing $x=2^3\sqrt{2}a$, $y=5\sqrt{5}a$ we have that

$$z^3 + 2^4(2^7a^2 + 10\sqrt{5}az + z^2)(40\sqrt{5}a + 5z) = 0 \quad (2.3)$$

has a double root. Since for its derivative we have

$$3z^2 + 2^4(10\sqrt{5}a + 2z)(40\sqrt{5}a + 5z) + 2^45(2^7a^2 + 10\sqrt{5}az + z^2) = 0$$

$$(15 + \frac{3}{16})z^2 + 180\sqrt{5}az + 2640a^2 = 0$$

$$\frac{a}{z} = \frac{-90\sqrt{5} \pm \sqrt{90^25 - 2640(15 + \frac{3}{16})}}{2640} = \sqrt{5} \frac{-30 \pm 3}{880}.$$

Thus $\frac{y}{z} = \frac{-25(30 \pm 3)}{880}$, $\frac{x}{z} = \frac{-8\sqrt{10}(30 \pm 3)}{880}$ and the point of tangency is one of the points $(-33 \cdot 8\sqrt{10}, -25 \cdot 33, 880)$, $(-27 \cdot 8\sqrt{10}, -25 \cdot 27, 880)$.

Upon substituting these values in the polynomial 2.3 we find that the correct choice is $(-24\sqrt{10}, -75, 80)$.

By symmetry the point $(24\sqrt{10}, -75, 80)$ is the tangency point of the line L_- .

Let us write

$$C = 4^2(8y+5z)x^2 + z(16y+9z)^2 = 0$$

and let us set $u=16y+9z$. We have:

$$C = zu^2 + 8x^2(u+z) = 0$$

In these coordinates the singular point of C is $(0, 0, 1)$, so the tangents at the singular point are given by:

$$8x^2 + u^2 = 0$$

whence they are complex and we have an isolated point.

In order to draw C , let's compute its flexes. Using the coordinates x , u , and z the Hessian matrix is:

$$\begin{pmatrix} 16(u+z) & 16x & 16x \\ 16x & 2z & 2u \\ 16x & 2u & 0 \end{pmatrix}$$

The Hessian curve is then given by the determinant of

$$\begin{pmatrix} (u+z) & 0 & x \\ 0 & z-2u & u \\ 8x & u & 0 \end{pmatrix}$$

which equals

$$-(u+z)u^2 - 8x^2(z-2u) = 0.$$

Eliminating $8x^2$ from the two equations we get

$$(u+z)^2 u^2 - zu^2(z-2u) = 0$$

so either $u = 0$, and this implies either $x=0$ (the singular point) or $z=0$ that gives the point $(1, 0, 0)$, or

$$(u+z)^2 - z(z-2u) = u^2 + 4uz = 0$$

that gives ($u \neq 0$) $u = -4z$, that is $z = -1$, $u = 4$, $y = \frac{13}{16}$, $x = \pm \sqrt{2/3}$.
For these points $\frac{x}{y} = \pm \sqrt{2/3} \frac{16}{13}$.




Figure 1: A basis of $\pi_1(\mathbb{P}^1 \setminus \{P_1, P_{-1}, P_+, P_-, P_0\})$.

3 Fundamental groups.

In this section we are going to describe the five steps leading to the determination of the orbifold fundamental group of the Persson's surfaces.

Step 1.

Let $\mathbb{F}_1$ be the blow up of $\mathbb{P}^2$ at the point $(0, 0, 1)$ and let Σ_∞ be the exceptional divisor.

We consider the fibre bundle $\mathbb{F}_1 \xrightarrow{f'} \mathbb{P}^1$ and its restriction f

$$\mathbb{F}_1 \setminus (C \cup Q \cup \Sigma_\infty \cup L_1 \cup L_{-1} \cup L_+ \cup L_- \cup L_0) = \tilde{\mathbb{F}}_1$$

$$f \downarrow$$

$$\mathbb{P}^1 \setminus \{P_1, P_{-1}, P_+, P_-, P_0\} = \mathbb{P}^1 \setminus \{5 \text{ pts.}\}.$$

f is again a fibre bundle and we have a corresponding homotopy exact sequence of fundamental groups

$$1 \longrightarrow \mathcal{F}_5 \longrightarrow \tilde{\Pi} \longrightarrow \mathcal{F}_4 \longrightarrow 1$$

where $\mathcal{F}_k$ denotes the free group with k generators and $\tilde{\Pi} = \pi_1(\tilde{\mathbb{F}}_1)$.

Here we choose a small positive real number ε and $x=\varepsilon$, $y=1$ as base point on $\mathbb{P}^1 \setminus \{5 \text{ pts.}\}$ and $x=\varepsilon$, $y=1$, $z=-4\sqrt{-1}$ as base point on $\mathbb{F}_1$.

We let $\delta_1, \dots, \delta_5$ be a natural geometric basis of the free group

$$\mathcal{F}_5 = \pi_1(f^{-1}(\text{base pt.})) = \pi_1(L_\varepsilon \setminus (C \cup Q \cup \Sigma_\infty))$$

where the five points $L_\varepsilon \cap C$, $L_\varepsilon \cap Q$ are ordered by lexicographic order on $\text{Re}(\frac{z}{y})$, $\text{Im}(\frac{z}{y})$.

$\mathcal{F}_4$ is generated by the five geometric paths γ'_i around the five critical values described in figure 1 and whose product is the identity.

For these elements we choose lifts to $\mathbb{F}_1$ using a C^∞ section of a tubular neighborhood of Σ_∞ meeting Σ_∞ just in the point ∞ ($y=0$) with intersection number equal to -1 .

Therefore such lifts give paths γ_i such that

$$\prod \gamma_i = \prod \delta_i$$

and more specifically

$$\gamma_+ \gamma_1 \gamma_0 \gamma_- \gamma_{-1} = \delta_1 \cdots \delta_5 = \gamma_{-1} \gamma_+ \gamma_1 \gamma_0 \gamma_-.$$

We have that, indeed, $\tilde{\Pi}$ occurs as a semidirect product described by the relations

$$\gamma_j^{-1} \delta_i \gamma_j = (\delta_i) \beta_j$$

where the β_j 's are suitable braids in

$$\begin{aligned} \mathfrak{B}_5 = \langle \sigma_1, \dots, \sigma_4 \mid & \sigma_i \sigma_j = \sigma_j \sigma_i \quad \forall 1 \leq i < j+1 \leq 5 \\ & \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1} \quad \forall 1 \leq i < 4 \rangle \end{aligned}$$

the braid group on 5 strings which acts on the right on the free group $\mathcal{F}_5$ by the formulae

$$\begin{aligned} (\delta_h) \sigma_k &= \delta_h \quad \text{if } h \neq k, k+1 \\ (\delta_k) \sigma_k &= \delta_{k+1} \\ (\delta_{k+1}) \sigma_k &= \delta_{k+1}^{-1} \delta_k \delta_{k+1}. \end{aligned}$$

The braids β_j are constructed by following the motion of the five points of the intersection of $f'^{-1}(P)$ with $C \cup Q$ while P goes along γ'_j .

With our choice of the γ 's we have, as the reader can easily verify,

$$\begin{aligned} \beta_0 &= \sigma_4^{12} \sigma_2^2 \\ \beta_1 &= \sigma_1^{-1} \sigma_2 \sigma_3 \sigma_1 \sigma_2^{-1} \sigma_1 \\ \beta_{-1} &= \sigma_4^{-6} \sigma_2^{-1} \beta_1 \sigma_2 \sigma_4^6 \\ \beta_+ &= \sigma_1^{-2} \sigma_2 \sigma_3 \sigma_4 \sigma_3^{-1} \sigma_2^{-1} \sigma_1^2 \\ \beta_- &= \sigma_4^{-6} \sigma_2^{-1} \beta_+ \sigma_2 \sigma_4^6. \end{aligned}$$

Step 2.

By taking $\sqrt{\frac{x-y}{x+y}}$ we have a new fibre bundle $\mathbb{F}_2 \xrightarrow{g'} \mathbb{P}^1$ obtained by base change.

Under this base change the inverse image Q' of the conic Q splits into two sections of g' which we will denote by Q'_1 and Q'_2 . Again, by restriction we have a fibre bundle g

$$\hat{\mathbb{F}}_2 = \mathbb{F}_2 \setminus (C' \cup Q'_1 \cup Q'_2 \cup \Sigma'_\infty \cup \{8 \text{ fibres}\}) \xrightarrow{g} \mathbb{P}^1 \setminus \{8 \text{ pts.}\}.$$

Correspondingly we get an exact sequence

$$1 \longrightarrow \mathcal{F}_5 = \langle \delta_1, \dots, \delta_5 \rangle \longrightarrow \hat{\Pi} \longrightarrow \mathcal{F}_7 = \langle \gamma_0, \gamma_-, \gamma_+, \bar{\gamma}_0, \bar{\gamma}_-, \bar{\gamma}_+, \gamma_1^2 \rangle \longrightarrow 1$$

where $\bar{\gamma}_i = \gamma_i^{\gamma_1} = \gamma_1 \gamma_i \gamma_1^{-1}$ and $\hat{\Pi} = \pi_1(\hat{\mathbb{H}}_2)$.

The fact that $\mathcal{F}_7$ has seven generators as above follows since the double cover of $\mathbb{P}^1 \setminus \{5 \text{ pts.}\}$ corresponds to the homomorphism $\mathcal{F}_4 \rightarrow \mathbf{Z}_2$ sending $\gamma'_1, \gamma'_{-1} \mapsto \bar{1}$, and $\gamma'_0, \gamma'_+, \gamma'_- \mapsto \bar{0}$.

If we want to keep track of the eight critical values, we can also use $(\gamma_{-1}^2)^{\gamma_1}$ as a generator. In fact

$$(\delta_1 \cdots \delta_5)^2 = (\gamma_+ \gamma_1 \gamma_0 \gamma_- \gamma_{-1})(\gamma_{-1} \gamma_+ \gamma_1 \gamma_0 \gamma_-)$$

thus

$$\gamma_+ \gamma_0^{\gamma_1} \gamma_-^{\gamma_1} (\gamma_{-1}^2)^{\gamma_1} \gamma_+^{\gamma_1} \gamma_1^2 \gamma_0 \gamma_- = (\delta_1 \cdots \delta_5)^2.$$

The geometric meaning of the above formula is related to the fact that $(\Sigma'_\infty)^2 = -2$, and more precisely to the fact that the new generators of $\mathcal{F}_7$ lie in a C^∞ section meeting Σ'_∞ in one point with intersection number (-2) , and not meeting the other curves.

A presentation of $\hat{\Pi}$ is thus given by

$$\begin{aligned} \langle \delta_1, \dots, \delta_5, \gamma_0, \gamma_-, \gamma_+, \bar{\gamma}_0, \bar{\gamma}_-, \bar{\gamma}_+, \Gamma = \gamma_1^{-2} \mid & \gamma_0^{-1} \delta_i \gamma_0 = (\delta_i) \beta_0 \\ & \vdots \\ & \bar{\gamma}_0^{-1} \delta_i \bar{\gamma}_0 = (\delta_i) \beta_1 \beta_0 \beta_1^{-1} \\ & \vdots \\ & \Gamma \delta_i \Gamma^{-1} = (\delta_i) \beta_1^2 \rangle \end{aligned}$$

Step 3.

The fundamental group

$$\Pi' = \pi_1(\mathbb{H}_2 \setminus (C' \cup Q'_1 \cup Q'_2 \cup \Sigma'_\infty \cup L'_1 \cup L'_{-1}))$$

is a quotient of $\hat{\Pi}$. The presentation of Π' is readily accomplished simply by introducing in the above presentation the further relations

$$\gamma_0 = \gamma_- = \gamma_+ = \bar{\gamma}_0 = \bar{\gamma}_- = \bar{\gamma}_+ = 1.$$

Then Π' is presented as

$$\begin{aligned} \langle \delta_1, \dots, \delta_5, \Gamma \mid & \delta_i = (\delta_i) \beta_0 \quad \delta_i = (\delta_i) \beta_- \quad \delta_i = (\delta_i) \beta_+ \\ & \delta_i = (\delta_i) \beta_1 \beta_0 \beta_1^{-1} \quad \delta_i = (\delta_i) \beta_1 \beta_- \beta_1^{-1} \end{aligned}$$

$$\delta_i = (\delta_i)\beta_1\beta_+\beta_1^{-1} \quad \Gamma\delta_i\Gamma^{-1} = (\delta_i)\beta_1^2 >$$

Remark: with the new relations we get, setting $\Gamma_{-1} = (\gamma_{-1}^2)^{\gamma_1}$,

$$\Gamma_{-1}\Gamma = (\delta_1 \cdots \delta_5)^2$$

Step 4'.

We denote by $X_2^\#$ the non singular part of the double cover X_2 of $\mathbb{F}_2$ (branched over $C' \cup Q'_1 \cup Q'_2 \cup \Sigma'_\infty$) and by $Z_2^\#$ the complement in $X_2^\#$ of L_1'', L_{-1}'' , the respective inverse images of L_1', L_{-1}' .

We finally let $Y_2^\#$ be the double cover of $\mathbb{F}_2 \setminus (C' \cup Q'_1 \cup Q'_2 \cup \Sigma'_\infty \cup L_1' \cup L_{-1}')$.

Thus $Y_2^\# \subset Z_2^\# \subset X_2^\#$.

Clearly $\pi_1(Y_2^\#) = \ker(\Pi' \longrightarrow \mathbf{Z}_2)$, where $\delta_i \mapsto \bar{1}$ and $\Gamma \mapsto \bar{0}$, is generated by Γ , $\sigma = \delta_1\Gamma\delta_1^{-1}$, $A_i = \delta_1\delta_i$ ($i=1, \dots, 5$) and $B_j = \delta_j\delta_1^{-1}$ ($j=2, \dots, 5$).

To find the relations we apply the Reidemeister-Shreier rewriting process to the relations R_α of Π' and to the relations $\delta_1 R_\alpha \delta_1^{-1}$.

Step 4''.

Clearly, $\pi_1(Y_2^\#)$ maps onto $\pi_1(Z_2^\#)$ surjectively with kernel normally generated by δ_1^2 , $\delta_i^2 = B_i A_i$ ($i=2, \dots, 5$) and $(\delta_1 \cdots \delta_5)^2$, thus $\pi_1(Z_2^\#)$ is generated by $A_2, \dots, A_5, \Gamma$ and has for relations the relations coming from the rewriting of R_α , $\delta_1 R_\alpha \delta_1^{-1}$, and the rewriting of $(\delta_1 \cdots \delta_5)^2 = 1$, i.e. $A_2 A_3^{-1} A_4 A_5^{-1} A_2^{-1} A_3 A_4^{-1} A_5 = 1$.

Remark: This relation says that the four generators $A_2, \dots, A_5$ are the generators of $\pi_1(\text{fibre}) = \pi_1(\text{genus 2 curve})$.

Step 5.

Let $m=k+1$ and consider $X_{2m}^\#$, the non singular part of the m -fold cyclic cover of X_2 totally branched over L_1'' and L_2'' .

To find a presentation of $X_{2m}^\#$ we first need a presentation of the kernel of the map $\pi_1(Z_2^\#) \longrightarrow \mathbf{Z}_m$ such that $A_i \mapsto \bar{0}$ and $\Gamma, \sigma \mapsto \bar{1}$, and then we add the relations $\Gamma^m = \sigma^m = 1$.

Applying the Reidemeister-Shreier method, we find that the kernel is generated by $\Gamma^m, \Gamma^i A_j \Gamma^{-i}$ for $i=1, \dots, m-1$ and $j=2, \dots, 5$, by $\Gamma^i \sigma \Gamma^{-i-1}$ for $i=1, \dots, m-2$ and $\Gamma^{m-1} \sigma$; it has for relations the rewriting in the new generators of the relations R_α'' of $\pi_1(Z_2^\#)$ and the rewriting of $\Gamma^i R_\alpha'' \Gamma^{-i}$ for $i=1, \dots, m-1$.

4 Calculations.

Step 3.

We have

$$\begin{aligned} \beta_0 &= \sigma_4^{12} \sigma_2^2 \\ \beta_1 &= \sigma_1^{-1} \sigma_2 \sigma_3 \sigma_1 \sigma_2^{-1} \sigma_1 \\ \beta_{-1} &= \sigma_4^{-6} \sigma_2^{-1} \beta_1 \sigma_2 \sigma_4^6 \end{aligned}$$

$$\begin{aligned}\beta_+ &= \sigma_1^{-2}\sigma_2\sigma_3\sigma_4\sigma_3^{-1}\sigma_2^{-1}\sigma_1^2 \\ \beta_- &= \sigma_4^{-6}\sigma_2^{-1}\beta_+\sigma_2\sigma_4^6\end{aligned}$$

The relations $\delta_i=(\delta_i)\beta_0$ are equivalent to the two relations

$$(\delta_4\delta_5)^6 = (\delta_5\delta_4)^6 \quad (4.1)$$

$$\delta_2\delta_3 = \delta_3\delta_2. \quad (4.2)$$

The relations $\delta_i=(\delta_i)\beta_+$ amount to

$$\delta_5 = \delta_2^{-1}\delta_1^{-1}\delta_2\delta_1\delta_2. \quad (4.3)$$

In fact, here and in the sequel, we use the following argument: β_+ is a conjugate $\sigma\sigma_4\sigma^{-1}$ of the braid σ_4 and the braid σ_4 yields the relation $\delta_4=\delta_5$. Therefore, if we set $\delta'_4=(\delta_4)\sigma^{-1}$, $\delta'_5=(\delta_5)\sigma^{-1}$, we get the relation $\delta'_4=\delta'_5$. By our particular choice of σ

$$\begin{aligned}\delta'_5 &= \delta_5 \\ \delta'_4 &= (\delta_4)\sigma_3^{-1}\sigma_2^{-1}\sigma_1^2 \\ &= (\delta_3)\sigma_2^{-1}\sigma_1^2 \\ &= (\delta_2)\sigma_1^2 \\ &= (\delta_2^{-1}\delta_1\delta_2)\sigma_1 \\ &= \delta_2^{-1}\delta_1^{-1}\delta_2\delta_1\delta_2\end{aligned}$$

Similarly, the relations $\delta_i=(\delta_i)\beta_-$ are equivalent to the relation

$$\delta_3^{-1}\delta_1^{-1}\delta_3\delta_1\delta_3 = (\delta_4\delta_5)^{-3}\delta_5(\delta_4\delta_5)^3. \quad (4.4)$$

We write down, for convenience of the reader, the action of the braid β_1^{-1} , since the new relations $\delta_i=(\delta_i)\beta_1\beta_j\beta_1^{-1}$ will be obtained from the relations equivalent to $\delta_i=(\delta_i)\beta_j$ simply by applying the automorphism β_1^{-1} .

$$\begin{aligned}(\delta_1)\beta_1^{-1} &= \delta_1\delta_2\delta_3\delta_2^{-1}\delta_1^{-1}\delta_2^{-1}\delta_1\delta_2\delta_4\delta_2^{-1}\delta_1^{-1}\delta_2\delta_1\delta_2\delta_3^{-1}\delta_2^{-1}\delta_1^{-1} \\ (\delta_2)\beta_1^{-1} &= \delta_1\delta_2\delta_3\delta_2^{-1}\delta_1^{-1} \\ (\delta_3)\beta_1^{-1} &= \delta_2^{-1}\delta_1\delta_2\delta_4^{-1}\delta_2^{-1}\delta_1^{-1}\delta_2\delta_1\delta_2\delta_4\delta_2^{-1}\delta_1^{-1}\delta_2 \\ (\delta_4)\beta_1^{-1} &= \delta_2^{-1}\delta_1\delta_2 \\ (\delta_5)\beta_1^{-1} &= \delta_5\end{aligned}$$

Thus, the relations $\delta_i=(\delta_i)\beta_1\beta_0\beta_1^{-1}$ are equivalent to the relations

$$(\delta_1\delta_2)^6 = (\delta_2\delta_1)^6 \quad (4.5)$$

$$\delta_5\delta_3\delta_5^{-1}\delta_4^{-1}\delta_5\delta_4 = \delta_4^{-1}\delta_5\delta_4\delta_5\delta_3\delta_5^{-1} \quad (4.6)$$

where we have used 4.3.

The relations $\delta_i = (\delta_i)\beta_1\beta_+\beta_1^{-1}$ are equivalent to the relation

$$\delta_5 = \delta_2^{-1}\delta_1\delta_2\delta_4^{-1}\delta_5\delta_3\delta_5^{-1}\delta_4\delta_2^{-1}\delta_1^{-1}\delta_2 \quad (4.7)$$

where we have used 4.3.

The relations $\delta_i = (\delta_i)\beta_1\beta_-\beta_1^{-1}$ pop up to the relation

$$\delta_3^{-1}\delta_1\delta_2\delta_1^{-1}\delta_3 = (\delta_4\delta_5)^{-2}\delta_5(\delta_4\delta_5)^2. \quad (4.8)$$

In fact, using 4.3 we have $(\delta_4\delta_5)\beta_1^{-1} = \delta_1\delta_2$ and so

$$((\delta_4\delta_5)^{-3}\delta_5(\delta_4\delta_5)^3)\beta_1^{-1} = (\delta_1\delta_2)^{-4}\delta_2(\delta_1\delta_2)^4$$

On the other side,

$$(\delta_1\delta_3)\beta_1^{-1} = \delta_1\delta_2\delta_3\delta_5^{-1}\delta_4\delta_5\delta_3^{-1}\delta_5^{-1}\delta_4^{-1}\delta_5\delta_4\delta_2^{-1}\delta_1^{-1}\delta_2$$

and using 4.6 and again 4.3

$$(\delta_1\delta_3)\beta_1^{-1} = \delta_1\delta_2\delta_3\delta_4\delta_5\delta_3^{-1}\delta_5^{-1}\delta_2^{-1}\delta_1^{-1}\delta_2 = \delta_1\delta_2\delta_3\delta_4\delta_5\delta_3^{-1}\delta_2^{-1}\delta_1^{-1}.$$

With the same method we have

$$\begin{aligned} ((\delta_1\delta_3)^{-1}\delta_3\delta_1\delta_3)\beta_1^{-1} &= \delta_1\delta_2\delta_3\delta_5^{-1}\delta_4^{-1}(\delta_3^{-1}\delta_5^{-1}\delta_4^{-1}\delta_5\delta_4\delta_5\delta_3)\delta_4\delta_5\delta_3^{-1}\delta_2^{-1}\delta_1^{-1} \\ &= \delta_1\delta_2\delta_3\delta_5^{-1}\delta_4^{-1}(\delta_5^{-1}\delta_4^{-1}\delta_5\delta_4\delta_5)\delta_4\delta_5\delta_3^{-1}\delta_2^{-1}\delta_1^{-1} \\ &= \delta_1\delta_2\delta_3(\delta_4\delta_5)^{-2}\delta_5(\delta_4\delta_5)^2\delta_3^{-1}\delta_2^{-1}\delta_1^{-1} \end{aligned}$$

So the relation is

$$\delta_3(\delta_4\delta_5)^{-2}\delta_5(\delta_4\delta_5)^2\delta_3^{-1} = (\delta_1\delta_2)^{-5}\delta_2(\delta_1\delta_2)^5 = \delta_1\delta_2\delta_1^{-1}$$

where we have used 4.5.

Finally we have to write the relations $\Gamma\delta_i\Gamma^{-1} = (\delta_i)\beta_1^2$, i.e.

$$\Gamma\delta_1\Gamma^{-1} = \Gamma\delta_2\Gamma^{-1}\delta_4^{-1}\delta_2^{-1}\delta_1\delta_2\delta_4\Gamma\delta_2^{-1}\Gamma^{-1} \quad (4.9)$$

$$\Gamma\delta_2\Gamma^{-1} = \delta_2^{-1}\delta_1\delta_2\delta_3^{-1}\delta_5\delta_3\delta_2^{-1}\delta_1^{-1}\delta_2 \quad (4.10)$$

$$\Gamma\delta_3\Gamma^{-1} = \delta_4^{-1}\delta_2^{-1}\delta_1^{-1}\delta_2\delta_4\delta_2^{-1}\delta_1\delta_2\delta_3^{-1}\delta_5^{-1}\delta_3\delta_5\delta_3\delta_2^{-1}\delta_1^{-1}\delta_2\delta_4^{-1}\delta_2^{-1}\delta_1\delta_2\delta_4 \quad (4.11)$$

$$\Gamma\delta_4\Gamma^{-1} = \delta_4^{-1}\delta_2^{-1}\delta_1^{-1}\delta_2\delta_4\delta_2^{-1}\delta_1\delta_2\delta_4 \quad (4.12)$$

$$\Gamma\delta_5\Gamma^{-1} = \delta_5 \quad (4.13)$$

Step 4.

We take as Shreier set for the left cosets of the kernel the set $\{S_0 = 1, S_1 = \delta_1\}$, so applying the Reidemeister-Shreier method we get the generators $\Delta = \delta_1^2$, Γ , $\sigma = \delta_1\Gamma\delta_1^{-1}$, $A_i = \delta_1\delta_i$ and $B_i = \delta_i\delta_1^{-1}$ for $i=2, 3, 4, 5$. For the relations we must

rewrite the relations 4.1,...,4.13 and their conjugate by δ_1 in terms of the new generators. The rewriting process goes as follows (cf. [MKS], pages 86-98):

$$\begin{aligned} S_0\delta_1 &= S_1 & S_1\delta_1 &= \Delta S_0 \\ S_0\delta_i &= B_i S_1 & S_1\delta_i &= A_i S_0 \quad \text{for } i=2, 3, 4, 5 \\ S_0\Gamma &= \Gamma S_0 & S_1\Gamma &= \sigma S_1 \end{aligned}$$

We want to show that it suffices to rewrite only the relations 4.1,...,4.13. Observe that all our relations can be written in the form $W\delta_i W^{-1} = \delta_k$ for a suitable word W . Assume that Γ doesn't appear in the relation and do the rewriting after moding out by the relations

$$\Delta = B_i A_i = 1. \quad (4.14)$$

Since $S_0\delta_i = A_i^{-1} S_1$ and also $S_0\delta_i^{-1} = A_i^{-1} S_1$, if we write $W = \prod_{\lambda=1}^h \delta_{j_\lambda}^{\pm 1}$, the rewriting of $W\delta_i W^{-1}\delta_k^{-1}$ is given by

$$A_{j_1}^{-1} A_{j_2} \cdots A_i^{\pm 1} \cdots A_{j_1}^{-1} A_k$$

(note that $A_1=1$). The rewriting of the same relation conjugated by δ_1 yields instead

$$A_{j_1} A_{j_2}^{-1} \cdots A_i^{\mp 1} \cdots A_{j_1} A_k^{-1}.$$

We get thus two relations of respective form $U A_k = 1$, $U^{-1} A_k^{-1} = 1$, which are obviously equivalent.

If instead Γ appears in the relation, we have one of the 4.9,...,4.13 which are of the form $\Gamma\delta_i\Gamma^{-1} = W\delta_i W^{-1}$ where we can in fact assume that Γ doesn't appear in the word W .

The rewriting of $\Gamma\delta_i\Gamma^{-1}W\delta_i^{-1}W^{-1}$ yields, again a relation of the form

$$\Gamma A_i^{-1} \sigma^{-1} U^{-1} = 1,$$

whereas the rewriting of the conjugate by δ_1 gives a relation

$$\sigma A_i \Gamma^{-1} U = 1,$$

which is an equivalent relation.

For convenience of notation we shall keep the generators $B_i = A_i^{-1}$.

To calculate $\pi_1(Z_2^\#)$ we must add the rewriting of $(\prod_{i=1}^5 \delta_i)^2 = 1$ which gives

$$A_2 B_3 A_4 B_5 B_2 A_3 B_4 A_5 = 1.$$

We have thus that $\pi_1(Z_2^\#)$ is generated by $A_2, A_3, A_4, A_5, \Gamma$ and σ and has the following set of relations

$$(B_4 A_5)^6 = (B_5 A_4)^3 \quad (4.15)$$

$$B_3A_2 = B_2A_3 \quad (4.16)$$

$$B_5 = B_2^3 \quad (4.17)$$

$$B_3^3 = (B_5A_4)^6 B_5 \quad (4.18)$$

$$A_2^{12} = 1 \quad (4.19)$$

$$B_5A_3B_5A_4B_5A_4 = B_4A_5B_4A_5B_3A_5 \quad (4.20)$$

$$B_5 = B_2^2A_4B_5A_3B_5A_4B_2^2 \quad (4.21)$$

$$B_3B_2B_3 = (B_5A_4)^4 B_5 \quad (4.22)$$

$$\sigma A_2^2 \Gamma^{-1} = A_4 B_2^2 A_4 \quad (4.23)$$

$$\Gamma B_2 \sigma^{-1} = B_2^2 A_3 B_5 A_3 B_2^2 \quad (4.24)$$

$$\Gamma B_3 \sigma^{-1} = B_4 A_2^2 B_4 A_2^2 B_3 A_5 B_3 A_5 B_3 A_2^2 B_4 A_2^2 B_4 \quad (4.25)$$

$$\Gamma B_4 \sigma^{-1} = B_4 A_2^2 B_4 A_2^2 B_4 \quad (4.26)$$

$$\Gamma B_5 \sigma^{-1} = B_5 \quad (4.27)$$

$$A_2 B_3 A_4 B_5 B_2 A_3 B_4 A_5 = 1 \quad (4.28)$$

where $B_i = A_i^{-1}$.

Let's reduce this presentation. Using 4.17 relation 4.21 becomes

$$B_4 A_2 B_4 = B_5 A_3 B_5 \quad (4.29)$$

and with this 4.20 becomes

$$B_2^4 = 1$$

which implies 4.19, and changes 4.17 into

$$B_5 = A_2.$$

Moreover, using 4.22 and the last equation, relation 4.18 gives

$$(A_2 A_4)^2 = A_3 A_2 B_3^2$$

and with this, using also 4.16, 4.22 becomes

$$B_3 B_2 = A_2 A_3 \quad (4.30)$$

thus transforming 4.29 into

$$B_3 = B_4 A_2 B_4$$

which allows us to delete the generator A_3 . Upon substituting the expressions of A_5 and A_3 into 4.28 and 4.30 we have

$$A_2 A_4 = A_4 A_2$$

$$A_4^4 = 1.$$

We can then see that the relations 4.16,...,4.28 are equivalent to the following

$$\begin{aligned} A_5 &= A_2^{-1} & A_3 &= A_2^{-1} A_4^2 \\ A_2^4 &= A_4^4 = 1 & A_2 A_4 &= A_4 A_2 \\ \sigma A_2^2 \Gamma^{-1} &= A_2^2 A_4^2 \\ \Gamma A_2^{-1} &= A_2^{-1} \sigma \\ \Gamma A_2 A_4^2 &= A_2 A_4^2 \sigma \\ \Gamma A_4^{-1} &= A_4 \sigma \\ \Gamma A_2 &= A_2 \sigma. \end{aligned}$$

Step 5.

We take as Shreier set for the left cosets of the kernel the set

$$\{R_i = \Gamma^i \mid i=0, 1, \dots, m-1\}$$

and we apply the Reidemeister-Shreier method.

The generators are $\hat{\Gamma} = \Gamma^m$, $A_{2,i} = \Gamma^i A_2 \Gamma^{-i}$, $A_{4,i} = \Gamma^i A_4 \Gamma^{-i}$ for $i=0, \dots, m-1$
 $\sigma_i = \Gamma^i \sigma \Gamma^{-(i+1)}$ for $i=0, \dots, m-2$ and $\sigma_{m-1} = \Gamma^{m-1} \sigma$.

For the rewriting process we have

$$\begin{aligned} R_i A_j &= A_{j,i} R_i & \text{for } j=2, 4 \ i=0, \dots, m-1 \\ R_i \Gamma &= R_{i+1} & \text{for } i=0, \dots, m-2 \\ R_{m-1} \Gamma &= \hat{\Gamma} R_0 \\ R_i \sigma &= \sigma_i R_{i+1} & \text{for } i=0, \dots, m-2 \\ R_{m-1} \sigma &= \sigma_{m-1} R_0. \end{aligned}$$

Thus, taking indices $i \pmod{m}$ and adding (as we must) the relation $\hat{\Gamma} = 1$, we obtain the relations

$$\begin{aligned} A_{2,i}^4 &= A_{4,i}^4 = 1 \\ A_{2,i} A_{4,i} &= A_{4,i} A_{2,i} \\ \sigma_i A_{2,i+1}^2 &= A_{2,i}^2 A_{4,i}^2 \\ A_{2,i+1}^{-1} &= A_{2,i}^{-1} \sigma_i \\ A_{2,i+1} A_{4,i+1}^2 &= A_{2,i} A_{4,i}^2 \sigma_i \\ A_{4,i+1}^{-1} &= A_{4,i} \sigma_i \\ A_{2,i+1} &= A_{2,i} \sigma_i. \end{aligned}$$

To simplify this presentation we write

$$\begin{aligned}
\sigma_i &= A_{2,i}^2 A_{4,i}^2 A_{2,i+1}^2 \\
&= A_{2,i} A_{2,i+1}^{-1} \\
&= A_{4,i}^2 A_{2,i}^{-1} A_{2,i+1} A_{4,i+1}^2 = A_{2,i}^{-1} A_{4,i}^2 A_{4,i+1}^2 A_{2,i+1} \\
&= A_{4,i}^{-1} A_{4,i+1}^{-1} \\
&= A_{2,i}^{-1} A_{2,i+1}
\end{aligned}$$

From the last and the second equations we get

$$A_{2,i}^2 = A_{2,i+1}^2 = A_{2,0}^2$$

and from the first one, remembering that $A_{2,i}$ commutes with $A_{4,i}$ and that $A_{2,0}^4 = 1$,

$$\sigma_i = A_{4,i}^2.$$

The fourth equation then gives

$$A_{4,i} = A_{4,i+1} = A_{4,0}$$

which makes the last and the third relations equivalent. These two cancellation relations enable us to delete all the generators σ_j and $A_{4,i}$ for $i=1, \dots, m-1$. We may rewrite the five relations above as

$$\begin{aligned}
\sigma_i &= A_{4,0}^2 \\
A_{4,i} &= A_{4,0} \\
A_{4,0}^2 &= A_{2,i}^{-1} A_{2,i+1} \\
A_{4,0}^2 &= A_{2,i} A_{2,i+1}^{-1}.
\end{aligned}$$

Clearly the last two equations are equivalent and give

$$A_{2,2i} = A_{2,0} \quad A_{2,2i+1} = A_{2,0} A_{4,0}^2. \quad (4.31)$$

Moreover, if we add the relation $\sigma^m = 1$, that in the generators of $\pi_1(X_{2m}^\#)$ reads out as $\sigma_0 \sigma_1 \cdots \sigma_{m-1} = 1$, we get $A_{4,0}^{2m} = 1$, i.e., if m is odd, $A_{4,0}^2 = 1$, while if m is even we have no new relations. Observe that this is in accordance with the fact that in 4.31 the index is cyclic mod(m).

Summing up, we have a commutative group with only two generators, namely $a = A_{2,0}$ and $b = A_{4,0}$, such that $a^4 = 1$ and $b^4 = 1$ if m is even, $b^2 = 1$ if m is odd, i.e.

$$\pi_1(X_{2k+2}^\#) = \mathbb{Z}_4 \times \mathbb{Z}_4$$

if k is odd and

$$\pi_1(X_{2k+2}^\#) = \mathbb{Z}_4 \times \mathbb{Z}_2$$

if k is even.

References

- [Ca] Catanese, F., “Generalized Kummer surfaces and differentiable structures on Noether-Horikawa surfaces, part I”, Proceedings of the Conference for Calabi’s 70th birthday, Symposia Mathematica, Cambridge University Press (1996).
- [Do] Donaldson, S., “Polynomial invariants for smooth four manifolds”, Topology **29** (1990), pp. 257–315.
- [F-S] Fintushel, R. and Stern R., “Rational blowdowns of smooth 4-manifolds”, preprint.
- [Fr] Freedman, M., “The topology of four dimensional manifolds”, J. Diff. Geom. **17** (1982), pp. 357–454.
- [Ho] Horikawa, E., “Algebraic surfaces of general type with small c_1^2 , I”, Ann. of Math. **104** (1976) pp. 357–387.
- [Kr] Kronheimer, P. B., “Instantons invariants and flat connections on the Kummer surfaces”, Duke Math. J. **64** (1991) pp. 229–241.
- [K-M] Kronheimer, P. B. and Mrowka, T., “Embedded surfaces and the structure of Donaldson’s polynomial invariants”, J. Diff. Geom. **41** (1995) pp. 573–734.
- [MKS] Magnus, Karrass, Solitar, “*Combinatorial Group Theory*”, Interscience, John Wiley and Sons (1966).
- [No] Noether, M., “Zur theorie des eindeutigen Entsprechens algebraischer Gebilde”, part 2, Math. Ann. Bd **8** (1875), pp. 495–533.
- [Per] Persson, U., “Horikawa surfaces with large Picard numbers”, Math. Ann. **259** (1982), pp. 287–312.
- [P-T] Pidstrigach, V. and Tyurin, A., “Localization of the Donaldson invariants along the Seiberg-Witten classes”, to appear in Izvestija of Russian Acad. of Science.
- [W] Witten, E., “Monopoles and four-manifolds”, Math. Res. Lett. **1** (1994), pp. 769–796.